arXiv:1206.2801v3 [gr-qc] 19 Apr 2013

Induced Gravity Cosmological Model with Non-positively Defined Higgs Potential

I. Ya. Aref'eva[1]*, N. V. Bulatov[2]†, R. V. Gorbachev[1]‡, S. Yu. Vernov[3]§

[1]Steklov Mathematical Institute, Russian Academy of Sciences,
Gubkina st. 8, 119991, Moscow, Russia
[2]Faculty of Physics,
Lomonosov Moscow State University, Leninskie Gory 1, 119991, Moscow, Russia,
[3]Skobeltsyn Institute of Nuclear Physics,
Lomonosov Moscow State University, Leninskie Gory 1, 119991, Moscow, Russia

Abstract

We study dynamics of a scalar field with the non-positively defined Higgs potential non-minimally coupled to gravity. In this model the inflationary stage of the Universe evolution could change into the quasi-cyclic stage of the Universe evolution with oscillation behaviour of the Hubble parameter from positive to negative values. Depending on the initial conditions the Hubble parameter can perform either one or several cycles its evolution.

1 Introduction

Models with scalar fields play a central role in current description of the evolution of the Universe at the early epoch [1, 2, 3, 4, 5, 6]. However, predictions of simplest inflationary models are in disagreement with the Planck2013 results [7, 8]. The models with the Ricci scalar multiplied by a function the scalar field are intensively studied in the inflationary cosmology [9, 10, 11, 12, 13, 14, 15, 16]. In the last years, the Higgs inflation has attracted a lot of attention [12, 13, 14, 15, 16].

There is also a consideration of theories with nonlocal scalar fields [17] inspired by the string field theory [18] as inflationary models [19, 20, 21]. Solutions with a bounce in these models have been considered in [22, 23] and perturbations have been studied in [24, 25]. In

---

*arefeva@mi.ras.ru
†nick_bulatov@mail.ru
‡rgorbachev@mi.ras.ru
§svernov@theory.sinp.msu.ru

one of these models the scalar field is the tachyon of the NSR fermion string and the model has the form of a nonlocal Higgs model. Due to the effect of stretching the potential [19, 20] nonlocality destroys the relation between the coupling constant in the potential, the mass term and the value of the vacuum energy (cosmological constant) and produces the effective Higgs potential with the negative extra cosmological term [26]. This effective Higgs potential is non-positively defined. It turns out that in the case of a minimal coupling non-positivity of a potential leads to the change of modes: from expansion to contraction [26]. In the cyclic model [27] the ekpyrotic phase is generated by a scalar field with a negative potential and one has a bounce from a phase of contraction to a phase of expansion, see [28] for discussion of the Planck2013 results support of this model.

In this paper, we consider the induced gravity model, which action includes the term proportional to the Ricci scalar multiplied by the square of the scalar field. The model incorporates only one scalar field, in other words, the scalar field which couples to the Ricci scalar is the same field whose potential drives inflation. We consider the non-positively defined potential, which is the Higgs potential minus a positive constant. The characteristic property of models with non-positively defined potential is the existence of the forbidden domain of the phase plane, which corresponds to non-real Hubble parameter. By numerical calculations we explore the case, when the forbidden domain is separated on the two parts and show by numerical calculations that the phase trajectories are attracting to the boundary of the forbidden domain, touch it and go to infinity. During this evolution the Hubble parameter oscillates from positive to negative values. This quasi-cyclic behaviors end up on a point of the boundary of the forbidden region and after this the evolution changes and completes by monotonically contracting stage. The phenomenon of a quasi-cyclic stage of the Universe evolution is a subject of our paper. Note that we got some results for an arbitrary non-positively defined potential as well.

The plan is the following. We start, Section 2, from a short remind of the action and equations for the induced gravity cosmological models. We consider dynamics in the models with an non-positively defined Higgs potential. We show that there are solutions that wind up on the forbidden region on the phase plane. In Section 3, we discuss the model in the Jordan and Einstein frames. In Section 4, we study the special case $\xi = -1/6$, in which some equations, considering in Section 2, have no sense. Finally Section 5 is devoted to the conclusion.

2 Classical dynamics in cosmological models with non-minimal coupling and non-positively defined potentials

Cosmological models in framework of a non-minimally coupled theory are actively studied [9, 10, 11, 29, 30, 31, 32, 33, 34, 35] (see also [36, 37] and references therein). Let us consider the the induced gravity models described by the following action [11, 33]:

$$S = \int d^4x\sqrt{-g}\left[\frac{1}{2}\xi\phi^2 R - \frac{1}{2}\partial_\mu\phi\partial^\mu\phi - V(\phi)\right], \tag{1}$$

where ξ is the non-minimal coupling constant.

We study the evolution of a homogeneous scalar field on a spatially flat Friedmann–Robertson–Walker (FRW) universe with the metric

$$ds^2 = -dt^2 + a^2(t)\left(dx_1^2 + dx_2^2 + dx_3^2\right).$$

The Einstein equations in the FRW metric look as follows:

$$H^2 = \frac{V(\phi)}{3\xi\phi^2} + \frac{1}{6\xi}\left(\frac{\dot{\phi}}{\phi}\right)^2 - 2H\frac{\dot{\phi}}{\phi}, \tag{2}$$

$$3H^2 + 2\dot{H} = -2\frac{\ddot{\phi}}{\phi} - 4H\frac{\dot{\phi}}{\phi} - \frac{4\xi+1}{2\xi}\left(\frac{\dot{\phi}}{\phi}\right)^2 + \frac{1}{\xi\phi^2}V(\phi), \tag{3}$$

where differentiation with respect to time t is denoted by a dot, and the Hubble parameter is the logarithmic derivative of the scale factor: $H = \dot{a}/a$.

The equation for the field got by the variation of the action over the field is

$$\ddot{\phi} + 3H\dot{\phi} + V' - 6\xi\phi\left(2H^2 + \dot{H}\right) = 0. \tag{4}$$

Combining equations (2)–(4), we obtain the following system of the first order differential equations

$$\dot{\phi} = \psi\,, \tag{5}$$

$$\dot{\psi} = -3H\psi - \frac{\psi^2}{\phi} + \frac{1}{(1+6\xi)\phi}[4V(\phi) - \phi V'(\phi)], \tag{6}$$

$$\dot{H} = \frac{4H\psi}{(1+6\xi)\phi} + \frac{V'(\phi)}{(1+6\xi)\phi} - \frac{12\xi}{1+6\xi}H^2 - \frac{1+2\xi}{2\xi(1+6\xi)}\left(\frac{\psi}{\phi}\right)^2. \tag{7}$$

We assume that $\xi \neq -1/6$. The case $\xi = -1/6$ will be considered separately in Section 4. All numerical calculations have been made for $\xi > 0$.

Equation (2) is the quadratic equation for H:

$$H^2 + 2H\frac{\dot{\phi}}{\phi} - \frac{V(\phi)}{3\xi\phi^2} - \frac{1}{6\xi}\left(\frac{\dot{\phi}}{\phi}\right)^2 = 0, \tag{8}$$

and has the following solutions

$$H_\pm = -\frac{\dot{\phi}}{\phi} \pm \sqrt{\left(1+\frac{1}{6\xi}\right)\left(\frac{\dot{\phi}}{\phi}\right)^2 + \frac{V(\phi)}{3\xi\phi^2}}. \tag{9}$$

The function H is a continuous function, so, if $V(\phi) > 0$ for all ϕ, then evolution of the Universe in such model is described either only H_- or only H_+. It depends on initial conditions. If $V(\phi)$ is not positively defined, then it is possible that a part of evolution is described by H_-, whereas another part — by H_+. The simplest way to get a non-positively defined potential from the known positively defined one is to subtract a positive constant.

Let us calculate a derivative of the expression $H+\frac{\psi}{\phi}$ over time. Using system (5)–(7), it is easy to show that

$$\frac{d}{dt}\left[H+\frac{\psi}{\phi}\right]=\frac{1}{2\xi(6\xi+1)\phi^2}\left(8\xi(V-3\xi\phi^2H^2)-(1+6\xi+24\xi^2)\psi^2-36\xi\left[\xi-\frac{1}{18}\right]\phi\psi H\right). \tag{10}$$

We can eliminate the potential V from this formula, using equation (2). Also, it is useful to write the Hubble parameter as

$$H=\ -\frac{\psi}{\phi}+Q.$$

Now equation (10) has the following form

$$\dot{Q}=-\frac{6\xi+1}{2\xi}\left(\frac{\psi}{\phi}\right)^2-\frac{\psi}{\phi}Q. \tag{11}$$

On the $(\phi,\dot{\phi})$ plane there is the boundary of the forbidden region. At any point of this boundary $Q=0$, so we get that always $\dot{Q}\leqslant 0$ and $\dot{Q}<0$ at $\psi\neq 0$. This means that if a solution for system (5)–(7) with H_+ touches this boundary at $\psi\neq 0$, then it change to H_-, solutions with H_- can not touch this boundary at $\psi\neq 0$.

From equations (2) and (3) we get the following equation, which form does not depend on the potential:

$$2\dot{H}=\ -2\frac{\ddot{\phi}}{\phi}+2H\frac{\dot{\phi}}{\phi}-\frac{2\xi+1}{\xi}\left(\frac{\dot{\phi}}{\phi}\right)^2. \tag{12}$$

Substituting the value of H on the boundary of the forbidden region, we get the following equation

$$\frac{6\xi+1}{\xi}\left(\frac{\psi}{\phi}\right)^2=0. \tag{13}$$

It means the boundary of the forbidden region is not a solution for system (5)–(7) at any values of parameters. Only trivial solutions with $\psi=0$ can belong to this boundary. The above-mentioned results are valid for any non-positively defined potential.

Models of this type in the case of the positively defined Higgs potential have been considered in a lot of works (see, for example, [12, 13, 14, 15]). In this paper, dynamics in the model with the non-positively defined Higgs potential

$$V_H(\phi)=\frac{\varepsilon}{4}\left(\phi^2-b^2\right)^2-\Lambda,\qquad \Lambda>0, \tag{14}$$

where b and Λ are constants, is studied.

In the case of the $V_H(\phi)$ potential, the boundary of the forbidden region ($Q=0$) has the following form

$$(1+6\xi)\dot{\phi}^2=2\Lambda-\frac{\varepsilon}{2}\left(\phi^2-b^2\right)^2. \tag{15}$$

Most attention will be paid to the case when $\Lambda < \varepsilon b^4/4$. In this case, the forbidden region on the $(\phi, \dot{\phi})$ plane consists of two separate parts.

Numerical calculations give the following phase trajectory for system (5)–(7). As we see there are two stages of evolution on the phase plane. The first stage corresponds to some kind of the cyclic Universe, then the phase trajectory reaches the boundary of the forbidden region and the second stage starts, during this stage the Hubble parameter rapidly decreases and the Universe contracts (See 1).



Figure 1: The solution of system (5)–(7) at $\Lambda = 0.05$. We choose $b = 1$, $\epsilon = 10$, $\xi = 10$. The initial conditions are $\phi_0 = 2$, $\psi_0 = 0$, H_0 is calculated by (9) with sing "+". On the left picture, the phase diagram is present. It is magenta at the corresponding value of $H = H_+ > 0$, blue at $H = H_+ < 0$, red at $H = H_- > 0$, and green at $H = H_- < 0$ brown. On this picture, brown dashed line corresponds to $H_+ = 0$, brown dashed line with long dashes corresponds to $H_- = 0$, black line is the boundary of the forbidden region. On the right picture, the Hubble parameter as function of the cosmic time is present. Brown color means that $H = H_+$, whereas $H = H_-$ is drown in dark green color.

The fact which gives the change of the stages is that when the phase trajectory reaches the boundary of the forbidden region the square root in the equation for the Hubble parameter gets the "minus" sign.

Note that the obtained behaviors of the solutions is essentially different from the behaviors of the solutions in the case $\Lambda = 0$. At $\Lambda = 0$ the forbidden region is absent and the point $(H = 0,\ \phi = b,\ \psi = 0)$ is an attractive fixed point and the Hubble parameter is positive at any point (see Fig 2).

The cyclic stage can be interpreted as a heating stage. To estimate the period of this reheating we can estimate a period of oscillations for a limit trajectory - a trajectory which coincides with a boundary of the forbidden region $(1 + 6\xi)\frac{\dot{\phi}^2}{2} = -V(\phi)$, we remind that this trajectory is not a solution for the system of equations under consideration. A solution to



Figure 2: The solution of system (5)–(7) at $\Lambda = 0$. We choose $b = 1$, $\epsilon = 10$, $\xi = 10$. The initial conditions are $\phi_0 = 1.5$, $\psi_0 = 0$, H_0 is calculated by (9) with sing "+". The Hubble parameter is always H_+.

equation $\ddot{\phi} = -\frac{1}{1+6\xi}V'(\phi)$ with negative energy $E = \Lambda' - \frac{\varepsilon' b^4}{4}$, where $\Lambda' = \frac{\Lambda}{1+6\xi}$, has the form (see, for example, [38])

$$\phi(t) = A\,\mathrm{dn}(\Omega t + c, k) \tag{16}$$

where $\mathrm{dn}(u, k)$ is the elliptic function of argument u and modulus k, A is the amplitude and c is the phase. The frequency Ω and the modulus k of the elliptic function are obtained from parameters ε', b of the equation and depend on the amplitude A:

$$A^2 \equiv b^2\left(1 + \sqrt{1 + \frac{4E}{\varepsilon' b^4}}\right) \tag{17}$$

$$\Omega^2 \equiv A^2\frac{\varepsilon'}{2} \tag{18}$$

$$k^2 \equiv 2\left(1 - \frac{b^2}{A^2}\right). \tag{19}$$

Parameter c is defined by the relation

$$\phi(0) = A\,\mathrm{dn}(c, k). \tag{20}$$

Now we estimate a period of function $\phi(t)$ (it will be of the same order as the period for the function H). Then we will also have an estimation for a period of H. Let us suppose the period has the standard form

$$T = \frac{2\pi}{\Omega} \tag{21}$$

and substitute Ω in the form (18). We get

$$T = 2\pi\sqrt{2}\sqrt{\frac{1+6\xi}{2\sqrt{\varepsilon\Lambda}+b^2\varepsilon}}. \tag{22}$$

Let us estimate the value of the period. For this purpose let us note that a non-minimal coupling turns the Planck mass into a dynamical quantity. The Planck mass is related to the vacuum expectation value of the potential b as follows [11]:

$$\frac{M_{pl}^2}{16\pi} = \frac{1}{2}\xi b^2 \tag{23}$$

This gives us an estimation for ξ

$$\xi = \frac{M_{pl}^2}{8\pi b^2}. \tag{24}$$

To estimate the value of the constant Λ let us note that we are considering the case when the forbidden area consists of two separate parts. It means that $\Lambda < \frac{\varepsilon b^4}{4}$, let us suppose that $\Lambda \sim \frac{\varepsilon b^4}{4}$. From $\Lambda \sim \frac{\varepsilon b^4}{4}$ it follows that

$$\sqrt{\varepsilon\Lambda} = \frac{\varepsilon b^2}{2}. \tag{25}$$

Then for a period it is the following estimation

$$T = \frac{3M_{pl}^2}{\sqrt{\varepsilon}b^2}. \tag{26}$$

For $b = 240GeV$ and $m_H = \sqrt{\varepsilon}b = 126GeV$ [12], [15] we get $\varepsilon \sim 1$ and for a period we have an estimation

$$T \sim 10^{14}GeV^{-1} \sim 10^{-11}s. \tag{27}$$

3 Action and equation of motion in the Einstein frame

Once a modified gravity theory is recast into its scalar-tensor presentation, it immediately follows that both the Jordan frame (where the scalar fields non-minimally couples to gravity) and the Einstein one (which has the same form as that of Einstein gravity with minimally coupled scalar fields) are available. These two frames are related by conformal transformation

$$g_{\mu\nu} = \Omega^2 g_{\mu\nu}^{(E)} \qquad \rightarrow \qquad g = \Omega^8 g^{(E)}\,, \tag{28}$$

where we denote the metric in the Jordan frame by $g_{\mu\nu}$, while the one in the Einstein frame is labeled as $g_{\mu\nu}^{(E)}$. In the following, we denote all quantities in the Einstein frame by adding a tag $^{(E)}$ to the corresponding ones in the other frame, in order to avoid confusion.

Let us consider the conformal transformation (28). Using $g^{\mu\nu} = \Omega^{-2}g^{\mu\nu(E)}$, one obtains the relationship between the Ricci scalars in the two frames [36]

$$R = \Omega^{-2}\left[R^{(E)} - 6\left(\Box^{(E)}\ln\Omega + g^{\mu\nu(E)}\nabla_\mu^{(E)}\ln\Omega\nabla_\nu^{(E)}\ln\Omega\right)\right]. \tag{29}$$

Inserting this into action (1), one identifies the conformal factor as

$$\Omega^{-2} = \kappa^2\xi\phi^2 \qquad \to \qquad \Omega = \frac{C}{\phi}, \tag{30}$$

where $C = 1/(\kappa\sqrt{\xi})$, where $\kappa^2 \equiv 8\pi/{M_{\rm Pl}}^2$, the Planck mass being $M_{\rm Pl} \simeq 1.22 \times 10^{19}$ GeV.

This results to the following action in the Einstein frame:

$$S = \int d^4x\sqrt{-g^{(E)}}\left\{\frac{1}{2\kappa^2}\left[R^{(E)} - \frac{12+C^2}{2\phi^2}g^{\mu\nu\,(E)}\partial_\mu\phi\partial_\nu\phi - \frac{C^4}{\phi^4}V(\phi)\right]\right\}. \tag{31}$$

Note that the term proportional to $\Box^{(E)}\ln(\phi)$ has been discarded as a total derivative. One can derive the equations of motion in the Einstein frame by varying this action.

As is well known [36], conformally flat metrics are mapped into each other. The FRW metric is conformally flat, so starting from the FRW metric in the Jordan frame we obtain the corresponding FRW metric in the Einstein one. This leads us to directly start from an FRW metric with cosmic time in the Einstein frame

$$ds^2 = \, - dt_E^2 + b_E^2(t_E)\delta_{ij}dx^idx^j\,, \tag{32}$$

where in dt_E and $a_E(t_E)$, the index E denotes the corresponding quantities in the Einstein frame. We get

$$dt_E = \frac{\phi}{C}dt, \qquad a_E = \frac{\phi}{C}a\,, \tag{33}$$

$$H_E \equiv \frac{d\log a_E}{dt_E} = \Omega\left(H - \frac{\dot{\Omega}}{\Omega}\right) = \frac{C}{\phi}\left(H + \frac{\psi}{\phi}\right) = \pm\frac{C}{\phi}\sqrt{\left(1+\frac{1}{6\xi}\right)\left(\frac{\dot{\phi}}{\phi}\right)^2 + \frac{V(\phi)}{3\xi\phi^2}}. \tag{34}$$

The Einstein frame is convenient for calculating the spectral index n_s for primordial perturbations [11]. It was shown that the equations (2) and (6) in this frame can be written as

$$\begin{aligned} H_E^2 &= \frac{\kappa^2}{3}\left[U(\varphi) + \frac{1}{2}\left(\frac{d\varphi}{dt_E}\right)^2\right], & (35)\\ \frac{d^2\varphi}{dt_E^2} + 3H_E\frac{d\varphi}{dt_E} &= -U'(\varphi), & (36)\end{aligned}$$

where the scalar field φ and the potential $U(\varphi)$ are

$$\frac{d\varphi}{d\phi} = \kappa^{-1}\sqrt{\frac{1+6\xi}{\xi\phi^2}}, \qquad U(\varphi) = \frac{V(\phi)}{(\kappa^2\xi\phi^2)^2}. \tag{37}$$

In slow-roll approximation the Einstein frame equations are

$$\begin{aligned} H_E^2 &\simeq \frac{\kappa^2}{3}U(\varphi),\\ 3H_E\frac{d\varphi}{dt_E} &\simeq -U'(\varphi).\end{aligned}$$

In other words, it has been assumed that the standard Einstein-frame "slow-roll" approximations may be made. This approximate equations in the Jordan frame look as

$$\begin{aligned}\left(H+\frac{\dot{\phi}}{\phi}\right)^2 &\simeq \frac{1}{3\xi\phi^2}V(\phi),\\ 3\left(H\dot{\phi}+\frac{\dot{\phi}^2}{\phi}\right) &\simeq \frac{1}{(1+6\xi)}\frac{1}{\phi}\left[4V(\phi)-\phi V'(\phi)\right].\end{aligned}$$

Yet the assumption, $U(\varphi)\gg\frac{1}{2}(d\varphi/dt_E\,)^2$, is equivalent to $V(\phi)\gg\frac{1}{2}\dot{\phi}^2(1+6\xi)$, and thus it remains consistent further to simplify the field equations during slow-roll as:

$$H^2\simeq\frac{V(\phi)}{3\xi\phi^2}\,, \tag{38}$$

$$3H\dot{\phi}\simeq\frac{1}{(1+6\xi)\phi}[4V(\phi)-\phi V'(\phi)]. \tag{39}$$

Using this approximation it is possible to obtain the values for the slow-parameters and the spectral index n_s as it has been done in [10]. In the case of non-positively defined potential the results will be qualitative the same. One can obtain both "chaotic" inflation and "new" inflation.

4 The special case $\xi=-1/6$

This case is of a special interest due to the fact that if $\xi=-1/6$ then equations (5)–(7) have no sense. To consider this case we put $\xi=-\frac{1}{6}$ into equations (2)–(4). These equations get the following form

$$H^2=-\frac{2V(\phi)}{\phi^2}-\left(\frac{\dot{\phi}}{\phi}\right)^2-2H\frac{\dot{\phi}}{\phi}, \tag{40}$$

$$3H^2+2\dot{H}=\left(\frac{\dot{\phi}}{\phi}\right)^2-2\frac{\ddot{\phi}}{\phi}-4H\frac{\dot{\phi}}{\phi}-\frac{6}{\phi^2}V(\phi), \tag{41}$$

$$\ddot{\phi}+3H\dot{\phi}+V'+\phi\left(2H^2+\dot{H}\right)=0. \tag{42}$$

Combining equations (40) and (41) we get

$$\dot{H}=2\left(\frac{\dot{\phi}}{\phi}\right)^2+2H\frac{\dot{\phi}}{\phi}+\frac{\ddot{\phi}}{\phi}. \tag{43}$$

After, from equation (42) we get the condition

$$4V(\phi)-\phi V'(\phi)=0. \tag{44}$$

This means that a solution with a nontrivial $\phi(t)$ exists if and only if the potential $V(\phi)=A\phi^4$. Constant A must be non-positive to satisfy the requirement that H is real. Equation (43) is an identity in this case. We get that for any function ϕ there is function H and they are solutions of equations.

5 Conclusion

In this paper, we have considered the with non-positively defined potential, which is the Higgs potential minus a positive constant. The characteristic property of models with non-positively defined potential is the existence of the forbidden domain of the phase plane, which corresponds to non-real Hubble parameter.

We have shown that behaviors of the Hubble parameter in with non-positively defined Higgs potential and Higgs potential are essentially different. For the Higgs potential we have attractive fixed points $\phi = \pm a$, which corresponds to minimums of the potentials. If the subtract a positive constant from the Higgs potential, then these fixed points appear in the forbidden domains. In the case of the non-positively defined Higgs potential the boundary of the forbidden region consists of two closed curves (if there is a special relation between a cosmological constant and parameters of the potential). We have explored this case and shown numerically that the phase trajectories are attracting to the boundary of the forbidden domain, touch it and go to infinity.

In this paper, we have considered the induced gravity model, which are very popular in cosmology [11, 30, 31]. Note that cosmological models with R, multiplied by $\xi\phi^2 + J$, where J is a constant, are considered as a real candidate to describe the inflationary scenario [12, 13, 15, 16], which is fully consistent with the Planck constraints [7]. We plan to consider this model in future investigations.

Acknowledgements. We would like to thank Dmitry Gorbunov for the helpful discussions. The work is partially supported by the RFBR grant 11-01-00894-a. The research of I.A., N.B., and R.G. is partially supported by grant NS − 4612.2012.1. N.B. and R.G. are also partially supported by grant the RFBR 12-01-31144. The research of S.V. is supported in part by the Russian Ministry of Education and Science under grant NSh-3920.2012.2.

References

[1] A.A. Starobinsky, *Relict Gravitation Radiation Spectrum and Initial State of the Universe* (In Russian), JETP Lett. **30** (1979) 682 [Pisma Zh. Eksp. Teor. Fiz. **30** (1979) 719];
A.A. Starobinsky, *A New Type of Isotropic Cosmological Models Without Singularity*, Phys. Lett. B **91** (1980) 99

[2] V.F. Mukhanov and G.V. Chibisov, *Quantum Fluctuation and Nonsingular Universe* (In Russian), JETP Lett. **33** (1981) 532 [Pisma Zh. Eksp. Teor. Fiz. **33** (190081) 549].

[3] A.H. Guth, *The Inflationary Universe: A Possible Solution to the Horizon and Flatness Problems*, Phys. Rev. D **23** (1981) 347

[4] A.D. Linde, *A New Inflationary Universe Scenario: A Possible Solution of the Horizon, Flatness, Homogeneity, Isotropy and Primordial Monopole Problems*, Phys. Lett. B **108** (1982) 389

[5] A. Albrecht and P.J. Steinhardt, *Cosmology for Grand Unified Theories with Radiatively Induced Symmetry Breaking*, Phys. Rev. Lett. **48** (1982) 1220

[6] A.D. Linde, *Chaotic Inflation*, Phys. Lett. B **129** (1983) 177

[7] P.A.R. Ade, *et. al.* [Planck Collaboration], *Planck 2013 results. XXII. Constraints on inflation*, arXiv:1303.5082

[8] P.A.R. Ade, *et. al.* [Planck Collaboration], *Planck 2013 results. XVI. Cosmological parameters*, arXiv:1303.5076;
P.A.R. Ade, *et. al.* [Planck Collaboration], *Planck 2013 Results. XXIV. Constraints on primordial non-Gaussianity*, arXiv:1303.5084

[9] B.L. Spokoiny, *Inflation And Generation Of Perturbations In Broken Symmetric Theory Of Gravity*, *Phys. Lett.* B **147** (1984) 39;
T. Futamase and K.-i. Maeda, *Chaotic Inflationary Scenario In Models Having Nonminimal Coupling With Curvature*, Phys. Rev. D **39** (1989) 399;
D.S. Salopek, J.R. Bond and J.M. Bardeen, *Designing Density Fluctuation Spectra in Inflation*, Phys. Rev. D **40** (1989) 1753;
R. Fakir and W.G. Unruh, *Improvement on cosmological chaotic inflation through nonminimal coupling*, *Phys. Rev.* D **41** (1990) 1783;
A. Tronconi and G. Venturi, *Quantum Back-Reaction in Scale Invariant Induced Gravity Inflation*, Phys. Rev. D **84** (2011) 063517, arXiv:1011.3958

[10] D.I. Kaiser, *Induced-gravity Inflation and the Density Perturbation Spectrum*, Phys. Lett. B **340** (1994) 23–28, arXiv:astro-ph/9405029

[11] D.I. Kaiser, *Primordial spectral indices from generalized Einstein theories*, Phys. Rev. D **52** (1995) 4295, arXiv:astro-ph/9408044

[12] F.L. Bezrukov and M. Shaposhnikov, *The Standard Model Higgs boson as the inflaton*, Phys. Lett. B **659** (2008) 703, arXiv:0710.3755

[13] F. Bezrukov, D. Gorbunov and M. Shaposhnikov, *On initial conditions for the Hot Big Bang*, JCAP **0906** (2009) 029, arXiv:0812.3622

[14] J. Garcia-Bellido, D.G. Figueroa, and J. Rubio, *Preheating in the Standard Model with the Higgs-Inflaton coupled to gravity*, Phys. Rev. D **79** (2009) 063531, arXiv:0812.4624

[15] A.O. Barvinsky, A.Y. Kamenshchik, C. Kiefer, A.A. Starobinsky, and C.F. Steinwachs, *Asymptotic freedom in inflationary cosmology with a non-minimally coupled Higgs field*, JCAP **0912** (2009) 003, arXiv:0904.1698;
A.O. Barvinsky, A.Y. Kamenshchik, C. Kiefer, A.A. Starobinsky, and C.F. Steinwachs, *Higgs boson, renormalization group, and cosmology*, Eur. Phys. J. C **72** (2012) 2219, arXiv:0910.1041

[16] A.O. Barvinsky, A.Y. Kamenshchik, C. Kiefer, and C.F. Steinwachs, *Tunneling cosmological state revisited: Origin of inflation with a non-minimally coupled Standard Model Higgs inflaton*, Phys. Rev. D **81** (2010) 043530, arXiv:0911.1408

[17] I.Ya. Aref'eva, *Nonlocal string tachyon as a model for cosmological dark energy*, AIP Conf. Proc. **826** (2006) 301, arXiv:astro-ph/0410443

[18] I.Ya. Arefeva, D.M. Belov, A.A. Giryavets, A.S. Koshelev and P.B. Medvedev, *Noncommutative field theories and (super)string field theories*, hep-th/0111208

[19] J.E. Lidsey, *Stretching the inflaton potential with kinetic energy*, Phys. Rev. D **76** (2007) 043511, arXiv:hep-th/0703007.

[20] N. Barnaby and J.M. Cline, *Large Nongaussianity from Nonlocal Inflation*, JCAP **0707** (2007) 017, arXiv:0704.3426

[21] I.Ya. Aref'eva and I.V. Volovich, *Cosmological Daemon*, JHEP **1108** (2011) 102, arXiv:1103.0273

[22] I.Ya. Aref'eva and L.V. Joukovskaya, *Time lumps in nonlocal stringy models and cosmological applications*, JHEP **0510** (2005) 087, hep-th/0504200

[23] I.Ya. Aref'eva, L.V. Joukovskaya and S.Yu. Vernov, *Bouncing and accelerating solutions in nonlocal stringy models*, JHEP **0707** (2007) 087, arXiv:hep-th/0701184;
I.Ya. Aref'eva, L.V. Joukovskaya, and S.Yu. Vernov, *Dynamics in nonlocal linear models in the Friedmann–Robertson–Walker metric*, J. Phys. A: Math. Theor. **41** (2008) 304003, arXiv:0711.1364

[24] A.S. Koshelev and S.Yu. Vernov, *Cosmological perturbations in SFT inspired non-local scalar field models*, Eur. Phys. J. C **72** (2012) 2198, [arXiv:0903.5176];
A.S. Koshelev and S.Yu. Vernov, *Analysis of scalar perturbations in cosmological models with a non-local scalar field*, Class. Quant. Grav. **28** (2011) 085019, [arXiv:1009.0746]

[25] F. Galli and A.S. Koshelev, *Perturbative stability of SFT-based cosmological models*, JCAP **1105** (2011) 012, arXiv:1011.5672

[26] I.Ya. Aref'eva, N.V. Bulatov and R.V. Gorbachev, *Friedmann cosmology with nonpositive-definite Higgs potentials*, Theor. Math. Phys. **173** (2012) 1466–1480, arXiv:1112.5951

[27] P. J. Steinhardt and N. Turok, *Cosmic Evolution in a Cyclic Universe*, Phys. Rev. D **65** (2002) 126003, hep-th/0111098

[28] J. -L. Lehners and P. J. Steinhardt, *Planck 2013 results support the simplest cyclic models*, arXiv:1304.3122 [astro-ph.CO].

[29] F. Cooper and G. Venturi, *Cosmology and broken scale invariance*, Phys. Rev. D **24** (1981) 3338

[30] A. Cerioni, F. Finelli, A. Tronconi and G. Venturi, *Inflation and Reheating in Induced Gravity*, Phys. Lett. B **681** (2009) 383–386, arXiv:0906.1902;
A. Cerioni, F. Finelli, A. Tronconi and G. Venturi, *Inflation and Reheating in Spontaneously Generated Gravity*, Phys. Rev. D **81** (2010) 123505, arXiv:1005.0935

[31] A.Y. Kamenshchik, A. Tronconi and G. Venturi, *Dynamical Dark Energy and Spontaneously Generated Gravity*, Phys. Lett. B **713** (2012) 358, arXiv:1204.2625

[32] J.L. Cervantes-Cota, R. de Putter, and E.V. Linder, *Induced Gravity and the Attractor Dynamics of Dark Energy/Dark Matter*, J. Cosmol. Astropart. Phys. **1012** (2010) 019, arXiv:1010.2237

[33] A.Yu. Kamenshchik, A. Tronconi, and G. Venturi, *Reconstruction of scalar potentials in induced gravity and cosmology*, Phys. Lett. B **702** (2011) 191–196, arXiv:1104.2125

[34] M. Sami, M. Shahalam, M. Skugoreva, and A. Toporensky, *Cosmological dynamics of non-minimally coupled scalar field system and its late time cosmic relevance*, Phys. Rev. D **86** (2012) 103532, arXiv:1207.6691

[35] A.Yu. Kamenshchik, A. Tronconi, G. Venturi, and S.Yu. Vernov, *Reconstruction of Scalar Potentials in Modified Gravity Models*, Phys. Rev. D **87** (2013) 063503, arXiv:1211.6272

[36] S. Capozziello and V. Faraoni, *Beyond Einstein Gravity: A Survey of Gravitational Theories for Cosmology and Astrophysics*, Fund. Theor. Phys. **170**, Springer, New York, 2011.

[37] S. Capozziello and M. De Laurentis, *Extended Theories of Gravity*, Phys. Rep. **509** (2011) 167, arXiv:1108.6266

[38] I.Ya. Arefeva, E.V. Piskovskiy and I.V. Volovich, *Rolling in the Higgs Model and Elliptic Functions*, Theor. Math. Phys. **172** (2012) 1001–1016, arXiv:1202.4395